Like Mixing Oil with Water

Assessing Contamination from Orphaned Oil Wells in the U.S.

Rolando Rabines

Env. Data Visualization | Tufts University

Abstract

This study investigates the relationship between methane concentrations in U.S. aquifers and their proximity to orphaned oil wells. Using spatial analysis, well data and methane contamination records were compared across different proximity thresholds. Results show a weak, negative correlation, with the highest methane levels found within 5 km. These findings highlight the environmental risks associated with oil and gas infrastructure and the need for remediation efforts.

Introduction


Orphaned oil wells pose environmental risks, including groundwater contamination with methane. This study aims to quantify this relationship by analyzing spatial links between oil wells and aquifers using methane concentrations as a key indicator.


Methods


Methane concentration data were available for a subset of water wells, limiting generalizability. Proximity was used as a proxy for contamination risk, without considering other factors like geological barriers or well depth. Data were sourced from the EPA, state agencies, Environmental Science & Technology publications, and USGS layers. Spatial analyses included linking orphaned wells with water wells via spatial join operations and nearest neighbor analysis. Visualizations featured maps and box-and-whisker plots showing methane concentrations across different proximity zones. Key limitations included data quality issues, temporal limitations, and geospatial resolution challenges.

Results

Spearman's rank correlation analysis between proximity (distance ranges) and methane concentrations yielded a very weak negative correlation coefficient ($\rho = -0.0137$) with a high p-value of 0.8222, indicating a minimal and statistically non-significant relationship between distance from orphaned oil wells and methane concentrations in water wells across the United States. Similarly, a Mann-Whitney U test comparing methane concentrations within a 5 km radius of orphaned oil wells with a random subset of water wells resulted in a test statistic of 702.5 and a p-value of 0.5789. This high p-value suggests no significant difference in methane concentrations between these two groups, reinforcing the conclusion that proximity to orphaned oil wells does not directly correlate with increased methane contamination in U.S. aquifers. The weak correlations may be due to the complex and multifaceted nature of methane contamination, influenced by factors such as geological barriers, local hydrology, and the presence of other pollutants, not solely proximity to orphaned oil wells.







References

1. Oil Industry Icon Set Licensed under CCBY 4.0
   Oil Industry Icon Set. (2024). Licensed under Creative Commons Attribution 4.0 International License. Retrieved from https://creativecommons.org/licenses/by/4.0/
2. Biden-Harris Administration Press Release
   The White House. (2023, November 29). Biden-Harris Administration invests $660 million to plug orphaned oil and gas wells. Retrieved https://www.doi.gov/pressreleases/biden-harris-administration-invests-660-million-states-plug-orphaned-oil-and-gas-wells
3. McMahon, P., Barlow, J., Engle, M., Belitz, K., Ging, P., Hunt, A., Jurgens, B., Kharaka, Y., Tollett, R., & Kresse, T. (2017). Methane and Benzene in Drinking-Water Wells Overlying the Eagle Ford, Fayetteville, and Haynesville Shale Hydrocarbon Production Areas. Environmental Science & Technology, 51(12), 6954-6963. https://doi.org/10.1021/acs.est.7b00746
4. DerelictWellMetadata
   Grove, C. A., Merrill, M. D. (2022, August 22). United States Documented Unplugged Orphaned Oil and Gas Well Dataset. U.S. Geological Survey. Retrieved from https://doi.org/10.5066/P91PJETI

Discussion

In the discussion, the results reveal a weak, negative correlation between the proximity of orphaned oil wells and methane concentrations in U.S. aquifers. The minimal correlation suggests that simply being close to orphaned oil wells may not directly translate to increased methane contamination risk. Certain factors likely contribute to this, including unmeasured variables such as geological barriers, well integrity, and the specific activities occurring at these sites. Additionally, limitations in the available methane concentration data for only a subset of water wells, coupled with a random sampling approach that may not fully capture regional variations in oil and gas activity, introduce potential biases. Measurement uncertainties due to variability in methane concentrations further dilute the observed relationship.

These findings highlight the complexity of contamination risks associated with orphaned oil wells and underscore the need for more comprehensive data and analyses to fully understand their environmental impact. Future research should explore additional factors influencing methane migration, including geological conditions, operational characteristics of oil wells, and temporal changes in contamination, to better assess the risk to groundwater resources.

Conclusion

Policymakers should prioritize remediation efforts in regions with high oil well densities to mitigate contamination risks. Further research should explore the persistence of methane contamination over time and its interactions with other pollutants.